

考 試 別：原住民族特考

等 別：四等考試

類 科 組：教育行政

科 目：教育測驗與統計概要

考試時間：1 小時 30 分

座號：_____

※注意：(一)可以使用電子計算器，須詳列解答過程。

(二)不必抄題，作答時請將試題題號及答案依照順序寫在試卷上，於本試題上作答者，不予計分。

(三)請以黑色鋼筆或原子筆在申論試卷上作答。

一、試解釋或比較下列成對的名詞：（每小題 5 分，共 20 分）

(一)認知測驗與情意測驗

(二)文字測驗與非文字測驗

(三)信度與效度

(四)團體測驗與個別測驗

二、T 分數與百分位數皆為常見之表示常模的方法，試回答有關此兩種常模的問題：

（每小題 15 分，共 30 分）

(一)試說明兩者有何異同？

(二)兩者在解釋測驗分數時之優缺點。

三、某校校長為了選擇運動服的顏色，於是從全校學生隨機抽取 100 人，施以顏色偏好問卷（參與者只能選一種他最喜歡的顏色），結果如下表 1 之觀察個數：（每小題 10 分，共 30 分）

表1：顏色偏好分布

	觀察個數	期望個數
白	19	20.0
紅	21	20.0
藍	27	20.0
綠	21	20.0
黃	12	20.0
總和	100	

(一)請你計算卡方（Chi-Square， χ^2 ）值。

(二)要檢定表 1 個數分布的適合度（goodness of fit），其卡方的自由度（degrees of freedom; df）是多少？

(三)若設顯著水準（ α ）為.05，請查表 2 可得 $.95\chi^2_{df}=?$ ，那麼能否推斷參與者對顏色偏好的分布是否均勻？

（請接第二頁）

考 試 別：原住民族特考

等 別：四等考試

類 科 組：教育行政

科 目：教育測驗與統計概要

表 2： χ^2 分配的自由度與百分點

df	1	2	5	10	20	30	50	70	80	90	95	98	99	99.9
1	.0002	.0006	.00393	.0158	.0642	.148	.455	1.074	1.642	2.706	3.841	5.412	6.635	10.827
2	.0201	.0404	.103	.211	.446	.713	1.386	2.408	3.219	4.605	5.991	7.824	9.210	13.815
3	.115	.185	.352	.584	1.005	1.424	2.366	3.665	4.642	6.251	7.815	9.837	11.341	16.268
4	.297	.429	.711	1.064	1.649	2.195	3.357	4.878	5.989	7.779	9.488	11.668	13.277	18.465
5	.554	.752	1.145	1.610	2.343	3.000	4.351	6.064	7.289	9.236	11.070	13.388	15.086	20.517
6	.872	1.134	1.635	2.204	3.070	3.828	5.348	7.231	8.558	10.645	12.592	15.033	16.812	22.457
7	1.239	1.564	2.167	2.833	3.822	4.671	6.346	8.383	9.803	12.017	14.067	16.622	18.475	24.322
8	1.646	2.032	2.733	3.490	4.594	5.527	7.344	9.524	11.030	13.362	15.507	18.168	20.090	26.125
9	2.088	2.532	3.325	4.168	5.380	6.393	8.343	10.656	12.242	14.684	16.919	19.679	21.666	27.877
10	2.558	3.059	3.940	4.865	6.179	7.267	9.342	11.781	13.442	15.987	18.307	21.161	23.209	29.588
11	3.053	3.609	4.575	5.578	6.989	8.148	10.341	12.899	14.631	17.275	19.675	22.618	24.725	31.264
12	3.571	4.178	5.226	6.304	7.807	9.034	11.340	14.011	15.812	18.549	21.026	24.054	26.217	32.909
13	4.107	4.765	5.892	7.042	8.634	9.926	12.340	15.119	16.985	19.812	22.362	25.472	27.688	34.528
14	4.660	5.368	6.571	7.790	9.467	10.821	13.339	16.222	18.151	21.064	23.685	26.873	29.141	36.123
15	5.229	5.985	7.261	8.547	10.307	11.721	14.339	17.322	19.311	22.307	24.996	28.259	30.578	37.697
16	5.812	6.614	7.962	9.312	11.152	12.624	15.338	18.418	20.465	23.542	26.296	29.633	32.000	39.252
17	6.408	7.255	8.672	10.085	12.002	13.531	16.338	19.511	21.615	24.769	27.587	30.995	33.409	40.790
18	7.015	7.906	9.390	10.865	12.857	14.440	17.338	20.601	22.760	25.989	28.869	32.346	34.805	42.312
19	7.633	8.567	10.117	11.651	13.716	15.352	18.338	21.689	23.900	27.204	30.144	33.687	36.191	43.820
20	8.260	9.237	10.851	12.443	14.578	16.266	19.337	22.775	25.038	28.412	31.410	35.020	37.566	45.315
21	8.897	9.915	11.591	13.240	15.445	17.182	20.337	23.858	26.171	29.615	32.671	36.343	38.932	46.797
22	9.542	10.600	12.338	14.041	16.314	18.101	21.337	24.939	27.301	30.813	33.924	37.659	40.289	48.268
23	10.196	11.293	13.091	14.848	17.187	19.021	22.337	26.018	28.429	32.007	35.172	38.968	41.638	49.728
24	10.856	11.992	13.848	15.659	18.062	19.943	23.337	27.096	29.553	33.196	36.415	40.270	42.980	51.179
25	11.524	12.697	14.611	16.473	18.940	20.867	24.337	28.172	30.675	34.382	37.652	41.566	44.314	52.620
26	12.198	13.409	15.379	17.292	19.820	21.792	25.336	29.246	31.795	35.563	38.885	42.856	45.642	54.052
27	12.879	14.125	16.151	18.114	20.703	22.719	26.336	30.319	32.912	36.741	40.113	44.140	46.963	55.476
28	13.565	14.847	16.928	18.939	21.588	23.647	27.336	31.391	34.027	37.916	41.337	45.419	48.278	56.893
29	14.256	15.574	17.708	19.768	22.475	24.577	28.336	32.461	35.139	39.087	42.557	46.693	49.588	58.302
30	14.953	16.306	18.493	20.599	23.364	25.508	29.336	33.530	36.250	40.256	43.773	47.962	50.892	59.703

Table C is adapted from Table IV of Fisher & Yates: *Statistical Tables for Biological, Agricultural and Medical Research*, published by Oliver & Boyd Ltd., Edinburgh, and by permission of the authors and publishers.

* If χ^2 is a chi-square variable with df greater than 30, then

$$z = \sqrt{2\chi^2} - \sqrt{2df - 1}$$

is very nearly normally distributed with mean 0 and standard deviation 1.

考 試 別：原住民族特考

等 別：四等考試

類 科 組：教育行政

科 目：教育測驗與統計概要

四、交通單位為宣導酒後不開車、開車不喝酒，於是想以科學證據勸導車輛駕駛人。宣導小組從有喝酒習慣的駕駛人中隨機抽樣 15 人，其實驗處理為先測量喝酒前的反應時間，接著測量這一組人喝酒後的反應時間。經初步統計分析後，結果如表 3 及表 4。
(每小題 10 分，共 20 分)

表3：成對樣本統計量

	平均數	個數	標準差	平均數的標準誤
喝酒前	30.33	15	7.267	1.876
喝酒後	34.33	15	5.690	1.469

表4：成對樣本t檢定

	成對變數差異					t值	自由度	顯著性 (雙尾)
	平均數	標準差	平均數的標準誤	差異的95%信賴區間				
				下界	上界			
成對	-4.000	2.976	.768	-5.648	-2.352	-5.205	14	.000

為了解釋上述實驗，請你完成下列問題，設顯著水準 (level of significance) α 為.05：

(一)陳述此實驗的統計假設 (虛無假設 vs.對立假設)。

(二)依表 4 的數據，你如何為此實驗做結論？(你可依表 4 資料或查表 5 做判斷)

考 試 別：原住民族特考

等 別：四等考試

類 科 組：教育行政

科 目：教育測驗與統計概要

表 5：t 分配的自由度與百分點

df	55	60	65	70	75	80	85	90	95	97.5	99	99.5	99.95
1	.158	.325	.510	.727	1.000	1.376	1.963	3.078	6.314	12.706	31.821	63.657	636.619
2	.142	.289	.445	.617	.816	1.061	1.386	1.886	2.920	4.303	6.965	9.925	31.598
3	.137	.277	.424	.584	.765	.978	1.250	1.638	2.353	3.182	4.541	5.841	12.941
4	.134	.271	.414	.569	.741	.941	1.190	1.533	2.132	2.776	3.747	4.604	8.610
5	.132	.267	.408	.559	.727	.920	1.156	1.476	2.015	2.571	3.365	4.032	6.859
6	.131	.265	.404	.553	.718	.906	1.134	1.440	1.943	2.447	3.143	3.707	5.959
7	.130	.263	.402	.549	.711	.896	1.119	1.415	1.895	2.365	2.998	3.499	5.405
8	.130	.262	.399	.546	.706	.889	1.108	1.397	1.860	2.306	2.896	3.355	5.041
9	.129	.261	.398	.543	.703	.883	1.100	1.383	1.833	2.262	2.821	3.250	4.781
10	.129	.260	.397	.542	.700	.879	1.093	1.372	1.812	2.228	2.764	3.169	4.587
11	.129	.260	.396	.540	.697	.876	1.088	1.363	1.796	2.201	2.718	3.106	4.437
12	.128	.259	.395	.539	.695	.873	1.083	1.356	1.782	2.179	2.681	3.055	4.318
13	.128	.259	.394	.538	.694	.870	1.079	1.350	1.771	2.160	2.650	3.012	4.221
14	.128	.258	.393	.537	.692	.868	1.076	1.345	1.761	2.145	2.624	2.977	4.140
15	.128	.258	.393	.536	.691	.866	1.074	1.341	1.753	2.131	2.602	2.947	4.073
16	.128	.258	.392	.535	.690	.865	1.071	1.337	1.746	2.120	2.583	2.921	4.015
17	.128	.257	.392	.534	.689	.863	1.069	1.333	1.740	2.110	2.567	2.898	3.965
18	.127	.257	.392	.534	.688	.862	1.067	1.330	1.734	2.101	2.552	2.878	3.922
19	.127	.257	.391	.533	.688	.861	1.066	1.328	1.729	2.093	2.539	2.861	3.883
20	.127	.257	.391	.533	.687	.860	1.064	1.325	1.725	2.086	2.528	2.845	3.850
21	.127	.257	.391	.532	.686	.859	1.063	1.323	1.721	2.080	2.518	2.831	3.819
22	.127	.256	.390	.532	.686	.858	1.061	1.321	1.717	2.074	2.508	2.819	3.792
23	.127	.256	.390	.532	.685	.858	1.060	1.319	1.714	2.069	2.500	2.807	3.767
24	.127	.256	.390	.531	.685	.857	1.059	1.318	1.711	2.064	2.492	2.797	3.745
25	.127	.256	.390	.531	.684	.856	1.058	1.316	1.708	2.060	2.485	2.787	3.725
26	.127	.256	.390	.531	.684	.856	1.058	1.315	1.706	2.056	2.479	2.779	3.707
27	.127	.256	.389	.531	.684	.855	1.057	1.314	1.703	2.052	2.473	2.771	3.690
28	.127	.256	.389	.530	.683	.855	1.056	1.313	1.701	2.048	2.467	2.763	3.674
29	.127	.256	.389	.530	.683	.854	1.055	1.311	1.699	2.045	2.462	2.756	3.659
30	.127	.256	.389	.530	.683	.854	1.055	1.310	1.697	2.042	2.457	2.750	3.646
40	.126	.255	.388	.529	.681	.851	1.050	1.303	1.684	2.021	2.423	2.704	3.551
60	.126	.254	.387	.527	.679	.848	1.046	1.296	1.671	2.000	2.390	2.660	3.460
120	.126	.254	.386	.526	.677	.845	1.041	1.289	1.658	1.980	2.358	2.617	3.373
∞	.126	.253	.385	.524	.674	.842	1.036	1.282	1.645	1.960	2.326	2.576	3.291

Table D is adapted from Table III of Fisher & Yates: *Statistical Tables for Biological, Agricultural and Medical Research*, published by Oliver & Boyd Ltd., Edinburgh, and by permission of the authors and publishers.

* The lower percentiles are related to the upper percentiles which are tabulated above by the equation $p_{f_n} = -_{1-p_{f_n}}$. Thus, the 10th percentile in the t -distribution with 15df equals the negative of the 90th percentile in the same distribution, i.e., $_{10}t_{15} = -1.341$.